

ANTI-PARKINSONISM DRUGS

BRAND NAMES

kineton	Dopar	Parda
lopa	Kemadrin	Parsidol
lopa	Larodopa	Phenoxe
	Levodopa	

PHENOTHIAZINES

BRAND NAMES

Proh	azine
	al
	an
	n

SCRIPTION DRUGS

TO CHOOSE

TO LOSE IT?

right to choose whether or
s under attack.

d that forcing individuals to
And those same politicians
ged from hospital, a leash
be dragged back again.

sked Mary Beth Valentine, Co-
dvocate Office, and David
o tell us what they think.

LL BE SERVED

l interest to consumers but
ome.

studies in
adequate studies in pre
avoid this drug during first 3
r guidance.

usually occurs
hyalactic reac-
ss, fever, sore
et.
ion, and faint-
e: Marked drowsiness, weakness, mus-
yness of mouth, blurred or double visio
Deep sleep progressing to coma, weal
w breathing, very low blood pressure, c
ded Use
Glossary).
inations While Taking
nts.

usually occurs
hyalactic reac-
ss, fever, sore
et.
ion, and faint-
e: Marked drowsiness, weakness, mus-
yness of mouth, blurred or double visio
Deep sleep progressing to coma, weal
w breathing, very low blood pressure, c
ded Use
Glossary).
inations While Taking
nts.

usually occurs
hyalactic reac-
ss, fever, sore
et.
ion, and faint-
e: Marked drowsiness, weakness, mus-
yness of mouth, blurred or double visio
Deep sleep progressing to coma, weal
w breathing, very low blood pressure, c
ded Use
Glossary).
inations While Taking
nts.

usually occurs
hyalactic reac-
ss, fever, sore
et.
ion, and faint-
e: Marked drowsiness, weakness, mus-
yness of mouth, blurred or double visio
Deep sleep progressing to coma, weal
w breathing, very low blood pressure, c
ded Use
Glossary).
inations While Taking
nts.

for Use by Those over 60 Years of Age.

Haloperidol

Possible Adverse Effects

IF ANY OF THE FOLLOWING
YOUR PHYSICIAN AS SOON

Mild Adverse Effects

DISCONTINUE DRUG AND NO

Chlorpromazi

You Need to Know for Safe Drug Use

serve the Following

restrictions. The liquid concentrate form of this

YOUR PHYSICIAN AS SOON AS POSSIBLE

Mild Adverse Effects

Allergic Reactions: Generalized itching, skin rashes

Other Reactions: Headache, dizziness, †drowsiness, †sluggishness, twitching, blurred vision.

Irregular pulse, heart palpitation.

Loss of appetite, nausea, †vomiting, †diarrhea

Serious Adverse Effects

Severe, involuntary, spasmodic movements, muscle rigidity, staggering gait.

Blackout spells, epileptic-like seizures.

Loss of bladder or rectal control.

CAUTION

1. This drug has a very narrow margin of safety. The dose to be effective is quite close to the level at which toxicity occurs. Careful dosage adjustments based on periodic blood level determinations are mandatory. Follow instructions on packaging and drug dosage.
2. Many over-the-counter (OTC) medications may interact unfavorably with this drug. Consult your physician before taking new medication of any kind.

Precautions for Use by Infants and Children

Initial dose should be low—not over 450 mg daily.

Increase dose at weekly intervals in steps of 150 mg.

Continue to increase until a blood level of 1.0 mEq/L is reached.

†May be an early sign of lithium toxicity, indicating the need for a blood level determination.

- propiomazine
- thiethylperazine
- thioridazine
- trifluoperazine
- triflupromazine
- trimeprazine



BARBITURATES
BRAND NAMES

- Mebaral
- Nembutal
- Sandoptal

TRANQUILIZERS, MAJOR
BRAND NAMES

- Miltown
- Serax
- Trancopal

THE RIGHT TO ARE WE GOING

The Mental Health Act guarantees our right to not to take medication. That right is being threatened.

Some politicians have been persuaded that staying on medication is good for them. They don't believe that even after we're discharged, we should be put on us so that we can't get better.

Let's talk about this threat. We've asked for the appointment of a Psychiatrist, Patient Advocate, M.P.P. and a former patient.

A FREE HOT MEAL WILL BE SERVED.

Note: This forum should be of special interest to mental health workers and therapists are also welcome.

- TRICYCLIC ANTIDEPRESSANTS**
BRAND NAMES
- Presamine
 - Sinequan
 - Surmontil
 - Tofranil
 - Vivacil

TRICYCLIC ANTIDEPRESSANTS

of no significance
Advisability of Use Depends on
Pregnancy Category:
Animal reproduction studies have shown that this drug may be harmful to the fetus. It is advisable to consult your physician.